

Bonner zoologische Beiträge	Band 55 (2006)	Heft 3/4	Seiten 203–222	Bonn, November 2007
-----------------------------	----------------	----------	----------------	---------------------

Biogeographic considerations of the Opisthobranchia (Mollusca: Gastropoda) fauna from the Brazilian littoral and nearby areas*

Francisco J. GARCÍA¹⁾, Marta DOMÍNGUEZ²⁾ & Jesús S. TRONCOSO²⁾

¹⁾Departamento de Sistemas Físicos, Químicos y Naturales; Facultad Ciencias Experimentales; Universidad Pablo de Olavide, Sevilla, Spain

²⁾Departamento de Ecología y Biología Animal, Facultad Ciencias del Mar, Universidad de Vigo, Vigo, Spain

*Paper presented to the 2nd International Workshop on Opisthobranchia, ZFMK, Bonn, Germany, September 20th to 22nd, 2006

Abstract. The aim of this paper is to examine the biogeographical distribution of the littoral and sub-littoral opisthobranch gastropods from Brazil and nearby areas. On the basis of published literature and personal data, a zoogeographic study was undertaken and the Brazilian region was compared with the Caribbean and Argentinean regions. The Brazilian littoral has been divided into six zones, based in the oceanographic features established by CASTRO & MIRANDA (1998). 466 species belonging to the orders Cephalaspidea, Anaspidea, Sacoglossa, Notaspidea and Nudibranchia from the Caribbean to the Argentinean region were considered in this study. The number of opisthobranch species is highest in Caribbean areas. Along the Brazilian littoral, South Brazilian Bight is the region with highest richness, while Orinoco-Amazapá and Amazon shelf are the areas with lower species numbers. The similarity analysis shows that some possible geographic barriers act to the distribution of the opisthobranchs. For each locality considered in the study, the percentage of species that extends northward is higher than southward.

Keywords. Brazil, Caribbean region, Argentinean region, biogeography, species list.

1. INTRODUCTION

Faunistic inventories give fundamental information for many basic and applied scientific disciplines, such as ecology and biogeography (STORK & SAMWAYS 1995). An essential tool to establish inventories is taxonomy, which supplies a reference system for the biodiversity (BISBY 1995). The level of faunistic and taxonomical knowledge from different regions varies considerably, which raises a problem for the accomplishment of biogeographic studies. Nevertheless, the elaboration of this type of studies offers insights into possible models of distribution of a taxonomic group throughout a more or less extensive area.

Opisthobranch gastropods are well represented in most marine habitats from equatorial to polar regions. Ernst and Eveline Marcus studied this group for more than thirty years in West Atlantic temperate and warmer waters. Recently, other authors have provided new contributions to the knowledge of the Brazilian opisthobranchs (TRONCOSO et al. 1998; GARCÍA et al. 2002; GARCÍA & TRONCOSO 2003, 2004; PADULA & ABSALÃO 2005; POLA et al. 2005; DOMÍNGUEZ et al. 2006 a, b; VALDÉS et al. 2006). The actual level of knowledge of the opisthobranchs from

Brazilian and Caribbean regions (i.e. MARCUS 1977; MARCUS & MARCUS 1967a; THOMPSON 1977, 1980; ESPINOSA & ORTEA 2001; ROSENBERG 2005; VALDÉS et al. 2006) permits some biogeographical considerations; however, future studies along some areas of Brazilian coast are necessary to obtain a more complete knowledge of this fauna. The work of Marcus and Marcus was focused around São Paulo and Rio de Janeiro areas. The necessity of more faunistic studies in Brazil is obvious when comparing the number of species cited along Brazilian coasts (280 species, after MARCUS 1977) with, for example, the Iberian Peninsula (523 species, after CERVERA et al. 2004).

Here, we present a study on the diversity of the Opisthobranchs along the Brazilian coasts, using as biogeographic areas the six zones defined by CASTRO & MIRANDA (1998). These Brazilian zones are also compared with Caribbean and Argentinean regions (i.e. MUNIAIN 1997; SCHRÖDL 1999; ROSENBERG 2005).

2. MATERIAL AND METHODS

2.1. Biogeographical areas

This research was conducted by comparing littoral and sub-littoral Opisthobranch fauna from the Brazilian shores with those from the Caribbean and Argentinean biogeographic regions. Species checklists were compiled by combining data from bibliographical sources and personal observations (included in the references). The total number of species considered for this study is 466, belonging to the orders Cephalaspidea, Anaspidea, Saeoglossa, Notaspidea and Nudibranchia (Table 1).

In this paper, we adopted the six oceanographic zones defined in CASTRO & MIRANDA (1998) for Brazil (Fig. 1). The geographical limits and features of these areas are shown in Table 2.



Fig. 1. Limits of the areas considered along Brazilian coasts, based on CASTRO & MIRANDA (1998) and nearby zones based on BRIGGS (1974). The number of species considered for each region is indicated.

Within the Caribbean region we have considered the provinces defined in BRIGGS (1974) (Fig. 1):

CC-CR, extends from Cape Canaveral to Cabo Romano

CR-CRo, extends from Cape Romano to Cape Rojo

CRo-O, extends from Cape Rojo to the mouth of the Orinoco River

O-A, extends from Orinoco River to Amapá

ABC, Includes Aruba, Bonaire and Curaçao

WI West Indies (WI)

Bermudas Is (BER).

In addition, we consider the Argentinean region (ARG) between the borders of Brazil to Uruguay up to 43-44°S (Chubut).

2.2. Community analysis and measurement of biodiversity

In order to examine diversity within the opisthobranch communities, data by region were subjected to a multivariate analysis using the Bray Curtis similarity measure and non-metric Multidimensional Scaling Ordination (MDS).

The Bray-Curtis index (BRAY & CURTIS 1957) was chosen because it does not consider double absences in its calculations. The results were then graphically described using dendrograms with the UPGMA (unweighted pair-group methods using arithmetic averages) aggregation algorithm (SNEATH & SOKAL 1973). The ordination analyses were carried out by means of an MDS (non metric multidimensional scaling program) based on the similarity matrix between stations.

For two different station groups a requirement is to identify which species account for the observed assemblage difference (CLARKE & GORLEY, 2001). The SIMPER routine was used to identify taxa that greatly contributed to differentiate station groups. The software used was P.R.I.M.E.R. (Plymouth Routines in Multivariate Ecological Research) version 5.2.8. for Windows.

3. RESULTS

466 species of opisthobranchs were found to occur in the Western Atlantic Ocean from Cape Canaveral to the Argentinean province (the Atlantic Magellanic region is not included). Table 3 shows the number of species by order or suborder for each zoogeographic area. The number of

opisthobranch species for each area varies remarkably. WI (West Indies) has the highest number of species (250 species), although, in general, the number is also high in the Caribbean areas Cro-O and CC-CR. Along the Brazilian zones, the number of species is lower and there is also a notable difference between the different zones. SBS is the zone with the highest number of species while AS is the zone with the lowest number (122 and 12 species, respectively). The number of species from the Argentinean region (ARG) is moderately low.

3.1. Faunal affinities

The cluster analysis using the six Brazilian zones shows a first division in which AS separate from the remaining zones (Fig. 2A). A second division separates SBS, and finally, the rest of the areas splits into two groups; a group including the north-eastern Brazilian zones EBS and NBS, and the other including ACR and SBB. A two-dimensional representation of the analysis MDS shows this same grouping pattern (Fig. 2B).






Fig. 2. Cluster classification (A) and MDS ordination (B) of the biogeographic areas based on the presence-absence of species in all areas included in this study. Black triangles, Brazilian regions; white triangles, Caribbean provinces; black circle, Argentinean region.

Including all geographical areas from Cape Canaveral and Bermudas to Chubut (Argentinean region), the cluster analysis and MDS show a first division in which Orinoco-Amapá (O-A), the Brazilian areas SBS and AS and the Argentinean region (ARG) separate from the remaining regions (Fig. 3). Those separate in two groups. One includes the north-eastern Brazilian zones EBS and NBS, while the other includes the remaining zones. The latter divides itself into two subgroups, one including the Brazilian regions SBB and ACR, and the other subgroup including all Caribbean provinces.

The SIMPER analysis identified four distinct groups. In group 1 (SBS, O-A, ARG, AS), the following taxa contribute to the similarity (up to 85%): Cephalaspidea (*Acteocina bidentata*, *Volvulella persimilis*, *Acteocina candei*, *Acteon pelecais*) and Notaspidea (*Pleurobranchaea inconspicua*). Group 2 (WI, CC-CR, CRo-O, CR-CRo, ABC, Ber) is mainly characterised by species of the or-

Fig. 2. Cluster classification (A) and MDS ordination (B) of the biogeographic areas based on the presence-absence of species in the areas from Brazil.

ders Cephalaspidea (*Volvulella*, *Bulla*, *Haminoea*, *Hydatina*, *Micromelo*), Notaspidea (*Umbraculum*, *Pleurobranchus*), Anaspidea (*Stylocheilus*, *Bursatella*, *Aplysia*, *Bosellia*), Sacoglossa (*Cylindrobulla*, *Oxynoe*, *Tridachia*, *Elysia*) Dendronotacea (*Scyllaea*) and Nudibranchia (*Spurilla*, *Chromodoris*). Group 3 (SBB, ACR) is characterised by the presence of species of the order Nudibranchia, mainly Doridina (*Dendrodoris krebsii*, *Cadlina rumia*, *Tyrinna evelinae*, *Diaulula greeleyi*, *Discodoris evelinae*, *Chromodoris clenchi*), Arminina (*Armina*) and Acolidina (*Spurilla neapolitana*, *Phidiana lynceus*, *Flabellina engeli*, *Glaucus atlanticus*). Group 4 (EBS, NBS) is determined (up to a cumulative 90%) by the presence of the orders Nudibranchia (*Doto divae*, *Diaulula greeleyi*), Notaspidea (*Pleurobranchaea inconspicua*, *Umbraculum umbraculum*, *Berthellina quadridens*), Anaspidea (*Phyllaplysia engeli*, *Stylocheilus striatus*, *Aplysia dactylomela*, *Aplysia parvula*, *Bursatella leachii*), Saccoglossa (*Cylindrobulla beauui*, *Oxynoe antillarum*, *Elysia tuca*), and Cephalaspidea (*Philine sagra*, *Hydatina vesicaria*, *Micromelo undatus*, *Haminoea elegans*, *Chelidonura petra*)

In terms of dissimilarity, the species *Spurilla neapolitana*, *Scyllaea pelagica*, *Pleurobranchus areolatus*, *Tridachia crispata*, *Aplysia dactylomela*, *Aplysia fasciata*, *Oxynoe antillarum* contribute greatly to differentiate groups 1 and 2. Group 1 differed from group 3 due to *Phidiana lynceus*, *Spurilla neapolitana*, *Flabellina engeli*, *Facelina coenda*, *Anteaeolidiella indica*, *Berghia benteva*, *Taringa telopia*, *Tyrinna evelinae*, *Siraius ilo*, *Jorinna spazzola*, *Okenia zoobotryon*, *Hallaxa aepae*, *Doris verrucosa*, *Discodoris evelinae*, *Chromodoris neona*, *Dendrodoris krebsii*, *Diaul-*

ula greeleyi, *Cadlina rumia*, *Chromodoris clenchi*, *Berthella agassizii*, *Berthella stellata*, *Aplysia dactylomela*, *Aplysia fasciata*, *Oxynoe antillarum*, *Ascobulla ulula*, *Navanax aenigmaticus*. Group 1 differed from group 4 due to *Doto divae*, *Diaulula greeleyi*, *Phyllaplysia engeli*, *Stylocheilus striatus*, *Berthellina quadridens*, *Aplysia dactylomela*, *Aplysia parvula*, *Oxynoe antillarum*, *Elysia tuca*, *Micromelo undatus*, *Chelidonura petra*, *Atys riseanus*. Differences between group 2 and group 4 were mainly due to *Tridachia crispata*, *Aplysia fasciata*, *Chelidonura petra*. The group 2 differed from group 3 due to *Facelina coenda*, *Berghia benteva*, *Scyllaea pelagica*, *Siraius ilo*, *Hallaxa aepae*, *Chromodoris neona*, *Tridachia crispata*, *Philine mera*, *Cylindrobulla beauui*, *Acteon pelecais*. And finally the species *Phidiana lynceus*, *Flabellina engeli*, *Glaucus atlanticus*, *Facelina coenda*, *Armina muelleri*, *Anteaeolidiella indica*, *Berghia benteva*, *Doto divae*, *Taringa telopia*, *Tyrinna evelinae*, *Siraius ilo*, *Okenia zoobotryon*, *Hallaxa aepae*, *Chromodoris neona*, *Dendrodoris krebsii*, *Cadlina rumia* explained most of the dissimilarity between groups 3 and 4.

Figure 4 shows the number and percentage of endemic species for each biogeographic area. Those species considered as endemic have been cited only at one zoogeographic region from the area of study. Geographic distribution along other biogeographic regions was not considered for this study. The level of endemism varies notably along the different zoogeographic areas. The highest value was found in the Argentinean region (68.6%). This high value is due to a southward distribution to the Magellan region of the fauna from this area. The Brazilian zones




Fig. 4. Number and percentage of endemic species for each faunistic area.

have percentages between 0 % in Amazon Shelf (AS) and East Brazilian Shelf (EBS), and 28.7 % in South Brazilian Bight (SBB). Among the Caribbean provinces, the percentage of endemism varies between 7.1 %, at Aruba, Bonaire and Curaçao (ABC), and 22 % in the West Indian region.

Table 4 shows the percentage of species from each region present in other localities. The percentage is increasing, when extended northward, and decreasing, when extended southward.

4. DISCUSSION

An aspect to be considered in this study is that the data on species distribution and the level of knowledge of the communities vary across the geographical regions. Thus, the results presented here must be considered as tentative.

The number of opisthobranch species between biogeographic areas varies remarkably being higher in the Caribbean: West Indies (WI: 250 species), Cape Rojo-Orinoco (CRo-O: 178) and Cape Canaveral-Cape Romano (CC-CR: 173). The south Brazilian Bight (SBB) is the zone in Brazil with the highest richness (122 species), followed by the north Brazilian Shelf (NBS: 75) and the Abrolhos-Campos Region (ACR: 62). The areas with the lowest species richness are Orinoco-Amapá (O-A) and Amazon Shelf (AS), both with 12 species, followed by South Brazilian Shelf (SBS) with 18 species. This could be related to oceanographic conditions, such as the effect of the Amazon River in the AS zone, and with a difference in sampling effort; the south Brazilian Bight is the zone where Ernst and Eveline Marcus did many of their collections.

The first division observed in the similarity analysis includes four geographical regions with features that seem to act as barriers to the distribution of opisthobranchs. The coasts from Orinoco-Amapá and Amazon Shelf are influenced by the Amazon and Orinoco rivers whose plumes spread north-westward for more than 1,000 km into the North Atlantic (CASTRO & MIRANDA 1998). These areas are characterised by soft bottom, turbid waters and freshwater runoff and they have been recognised as barriers to the dispersal of corals (COX & MOORE 2000), rocky shore gastropods (VERMEIJ 1978) and shallow water reef fishes (GILBERT 1972). The scarce opisthobranch fauna from Orinoco-Amapá and Amazon Shelf and the composition of species present in these areas are likely related to environmental characteristics of the region. Species present in these areas are generally cephalaspideans and arminacean, which frequently live in sand or mud.

A similar situation is found on the South Brazilian Shelf, where the low species numbers may be related to the effect of the Patos Lagoon River plume with an annual mean discharge of about $2000 \text{ m}^3 \text{ s}^{-1}$ (MARQUES et al. 2006). This area is influenced during the winter by Subantarctic water.

The coastal area of the Argentinean biogeographic province is a transition zone characterised by processes of mixing and instability of the water masses. This province includes geographical features like the Rio de la Plata. The estuaries influence the primary and secondary production in the area and consequently, the distribution of species. Moreover, there is an interaction off the coast between the Malvinas current flowing on the slope from the south with cold Subantarctic waters rich in nutrients and the Brazilian current, with temperatures higher than 20°C and salinity over 36.0 ppt. This determines the presence of eurythermal and euryhaline species (BOSCHI 2000). The fauna of opisthobranchs is formed mainly by nudibranchs, which have their northern distribution limit at the border between the Argentinean region and the South Brazilian Shelf, extending southwards to Subantarctic regions.

Cluster analyses show two Brazilian groups, composed of NBS-EBS and ACR-SBB. In general, these groups coincide geographically with those indicated by FLOETER et al. (2001) to the reef-fish fauna of the Brazilian coast. These authors considered several regions like the South and South-eastern coastal reefs, from the Guarapari islands to Santa Catarina (areas included in our analysis as the group ACR-SBB), and the North-eastern coast, extending from the Manuel Luis reefs to Abrolhos Archipelago (areas included in our analysis as the group NBS-EBS).

The southern and south-eastern coastal reefs, cited by FLOETER et al. (2001) show lower mean annual water temperature, relatively higher primary production and a large shelf width. The reef-fish fauna living in this area appears to be the richest of Brazil, due to the mix of tropical and subtropical elements. The area is subjected to a relatively intense seasonal upwelling promoted by the South Atlantic Central Water, bringing low-temperature ($<18^\circ \text{C}$) and nutrient-rich waters close to the coastline (EKAU & KNOPPERS 1999). FLOETER et al. (2001) stated that a considerable number of Caribbean reef fishes found in this region are absent from the north-eastern sites. We found a similar pattern in opisthobranchs. The cluster analysis groups ACR-SBB closer to the Caribbean provinces than to other Brazilian zones. On the other hand, ACR and SBB are the Brazilian areas with the highest richness in opisthobranchs. This could be related with environmental features (as is observed for reef-fishes; FLOETER et al. 2001). Nevertheless, in addition to environmental factors, differences

in the richness of opisthobranchs for each area may depend upon other factors, such as discrepancies in sampling effort. The fauna from Brazil is better known in the South Brazilian Bight and Abrolhos-Campos zone, where Evelyne and Ernst Marcus conducted research for over 30 years.

The north-eastern region as is described by FLOETER et al. (2001) for fishes, which nearly overlap with the group NBS-EBS, is characterised by its relatively warm waters, a weak seasonal signal and a small vertical temperature gradient; the circulation is influenced northward by the North Brazilian Current, and southward by the Brazilian Current (CASTRO & MIRANDA 1998). The reef formation consists of coralline algal crusts over a rocky substrate, hermatypic and fire corals, as well as sponges (FLOETER et al. 2001). The narrow and open shelf is an oligotrophic system almost entirely covered by carbonate sediments due to little freshwater input and the coast is influenced by the South Equatorial Current (KNOPPERS et al. 1999). In these areas herbivorous Saeoglossa and Anaspidea are more abundant. This trend of an increase in abundance in herbivores towards the tropical zone was previously observed in the Brazilian reef fishes (FERREIRA et al. 2004).

With regard to the distribution of opisthobranch species, it can be noted that for each locality considered in this study, the percentage of species that extend northward is higher than southward. Several authors discussed the Southern Caribbean as a centre of origin of species (see BRIGGS 2006) from which the species have been penetrating northward into Florida and Bermudas and southward into Brazilian waters (ROCHA 2003). Concerning Opisthobranchia, more intensive faunistic studies along the South American Atlantic coast are needed, before this hypothesis can be applied to understand opisthobranch diversity in that region.

Acknowledgments. This paper has been partially supported by the projects PHB2002-0045-PC and CGL2004-20366-E/BOS, of the Ministerio de Educación y Ciencia (MEC, Spain).

REFERENCES

- BISBY, F. A. 1995. Characterization of biodiversity. Pp. 21–106 in: HEYWOOD, V. H. (ed.) Global biodiversity assessment. Cambridge University Press. UCA.
- BOSCHI, E. E. 2000. Species of Decapod Crustaceans and their distribution in the American marine zoogeographic provinces. *Revista de Investigación y Desarrollo Pesquero* **13**: 7–136.
- BRAY R. J. & CURTIS, J. I. 1957. An ordination of the upland forest communities of Southern Wisconsin. *Ecological Monographs* **27**: 325–349.
- BRIGGS, J. C. 1974. *Marine Zoogeography*. New York: MacGraw-Hill.
- BRIGGS, J. C. 2006. Proximate sources of marine biodiversity. *Journal of Biogeography* **33**: 1–10.
- CASTRO, B. M. & MIRANDA, L. B. 1998. Physical Oceanography of the western Atlantic continental shelf located between 4°N and 34°S. *The Sea* **11**: 209–251.
- CERVERA, J. L.; CALADO, G.; GAVAIA, C.; MALAQUIAS, M. A. E.; TEMPLADO, J.; BALLESTEROS, M.; GARCÍA-GÓMEZ, J. C. & MEGINA, C. 2004. An annotated and updated checklist of the opisthobranchs (Mollusca: Gastropoda) from Spain and Portugal (including islands and archipelago). *Boletín Instituto Español de Oceanografía* **20**: 1–122.
- CLARKE, K. R. & GORLEY, R. N. 2001. *PRIMER (Plymouth Routines In Multivariate Ecological Research) v5: User manual/tutorial*: 1–91. (PRIMER-E Ltd, Plymouth).
- COX, C. B. & MOORE, P. D. 2000. *Biogeography. An ecological and evolutionary approach*. Blackwell Science, Oxford. Pp. 298.
- DOMÍNGUEZ, M., GARCÍA, F. J. & TRONCOSO, J. S. 2006 a. Some aspects of the family Chromodorididae (Opisthobranchia: Nudibranchia) from Brazil, with description of a new species. *Scientia Marina* **70** (4): 621–634.
- DOMÍNGUEZ, M., GARCÍA, F. J. & TRONCOSO, J. S. 2006 b. A new species of *Hoplodoris* Bergh, 1880, (Gastropoda: Opisthobranchia: Nudibranchia) from the Atlantic Ocean. *The Nautilus* **120** (4): 150–155.
- EDMUNDS, M. 1964. Eolid mollusca from Jamaica, with descriptions of two new genera and three new species. *Bulletin of Marine Science of the Gulf and Caribbean* **14**(1): 1–32.
- EDMUNDS, M. & JUST, H. 1983. Eolid nudibranchiate mollusca from Barbados. *Journal of Molluscan Studies* **49**: 185–203.
- EDMUNDS, M. & JUST, H. 1985. Dorid, Dendronotid and Arminid Nudibranchiate Mollusca from Barbados. *Journal of Molluscan Studies* **51**: 52–63.
- EKAU, W. & KNOPPERS, B. 1999. An introduction to the pelagic system of the north-east and east Brazilian shelf. *Archives of Fishery Marine Research* **47**: 113–132.
- ESPINOSA, J. & ORTEA, J. 2001. *Moluscos del Mar Caribe de Costa Rica: desde Cahuita hasta Gandoca*. *Avicennia Suplemento 4*: 1–77.
- FERREIRA, C. E. L.; FLOETER, S. R.; GASPARINI, J. L.; FERREIRA, B. P. & JOYEUX, J. C. 2004. Trophic structure patterns of Brazilian reef fishes: a latitudinal comparison. *Journal of Biogeography* **31**: 1093–1106.
- FLOETER, S. R.; GUIMARAES, R. Z. P.; ROCHA, L. A.; FERREIRA, C. E. L.; RANGEL, C. A. & GASPARINI, J. L. 2001. Geographic variation in reef-fish assemblages along the Brazilian coast. *Global Ecology and Biogeography* **10**: 423–433.
- GARCÍA, F. J. & TRONCOSO, J. S. 2003. Two unknown species of Mollusca Gastropoda from the Archipelago Fernando de Noronha (Brazil), with description of a new species belonging to the genus *Phidiana* Gray, 1850 and a new record of *Dendrodoris senegalensis* Bouchet, 1975. *Scientia Marina* **67** (2): 159–166.
- GARCÍA, F. J. & TRONCOSO, J. S. 2004. A new species of the genus *Anetarca* Gosliner, 1991 (Gastropoda: Opisthobranchia: Facelinidae) from the western Atlantic Ocean. *The Nautilus* **118** (4): 139–143.
- GARCÍA, F. J., TRONCOSO, J. S. & DOMÍNGUEZ, M. 2002. New data on the benthic Opisthobranch Molluscs from the Archipelago of Fernando de Noronha (Brazil), with description of a new species of Aegires Lovén. 1844. *Iberus* **20** (2): 45–56.
- GILBERT, C. R. 1972. Characteristics of the Western Atlantic Reef-fish Fauna. *Quarterly Journal of the Florida Academy of Sciences* **35**: 130–144.
- KNOPPERS, B.; EKAU, W. & FIGUEIREDO, A. G. 1999. The coast and shelf of east and northeast Brazil and material transport. *Geo-Marine Letters* **19**: 171–178.

- MARCUS, ER. 1955. Opisthobranchia from Brazil (1). Boletim de Universidade de São Paulo Faculdade de Filosofia, Ciências e Letras (Zoologia) **20**: 89–262.
- MARCUS, ER. 1957. On Opisthobranchia from Brazil (2). Journal of the Linnean Society of London **43** (292): 390–486.
- MARCUS, ER. 1958a. On western Atlantic opisthobranchiate gastropods. American Museum Novitates **1906**: 1–82.
- MARCUS, ER. 1958b. Notes on Opisthobranchia. Boletim do Instituto Oceanográfico, São Paulo **7** (1–2): 31–78.
- MARCUS, ER. 1961. Opisthobranchia from North Carolina. Journal of the Elisha Mitchell Scientific Society **77** (1): 141–151.
- MARCUS, EV. 1970a. Opisthobranchs from northern Brazil. Bulletin of Marine Science **20**: 922–951.
- MARCUS, EV. 1970b. On some Opisthobranchs from Cananéia, Brazil. Boletim Zoologia e Biologia Marinha, Universidade de São Paulo **27**: 207–219.
- MARCUS, EV. 1972. On some opisthobranchs from Florida. Bulletin of Marine Science **22**: 284–308.
- MARCUS, EV. 1972b. On the Anaspeidea (Gastropoda: Opisthobranchia) of the warm waters of the western Atlantic. Bulletin of Marine Sciences **22** (4): 841–874.
- MARCUS, EV. 1976a. On *Kentrodonis* and *Jorunna* (Gastropoda, Opisthobranchia). Boletim de Zoologia, Universidade de São Paulo **1**: 11–68.
- MARCUS, EV. 1976b. Opisthobranchia von Santa Marta, Columbia. Studies on Neotropical Fauna and Environment **11**: 119–150.
- MARCUS, EV. 1977. An annotated check list of the Western Atlantic warm water Opisthobranchs. Journal of Molluscan Studies **Supplement 4**: 1–22.
- MARCUS, EV. 1980. Review of Western Atlantic Elysiidae with a description of a new Elysia species. Bulletin of Marine Science **30** (1): 5–79.
- MARCUS, EV. 1983. The Western Atlantic Tritoniidae. Boletim de Zoologia, Universidade de São Paulo **6**: 177–214.
- MARCUS, EV. 1984. The Western Atlantic warm water Notaspidea (Gastropoda, Opisthobranchia), Parte 2. Boletim Zoologia Universidade São Paulo **8**: 43–76.
- MARCUS, EV. & HUGHES, H. P. I. 1974a. Opisthobranch mollusks from Barbados. Bulletin of Marine Science **24**: 498–532.
- MARCUS, EV. & HUGHES, H. P. I. 1974b. Opisthobranchs from American Atlantic warm waters. Bulletin of Marine Science of the Gulf and Caribbean **10**: 129–203.
- MARCUS, EV. & MARCUS, ER. 1955 (1957). Sea-hare and side-gilled slugs from Brazil. Boletim do Instituto Oceanográfico São Paulo **6**: 3–48.
- MARCUS, EV. & MARCUS, ER. 1958. Some opisthobranchs from the Northwestern Gulf of Mexico. Institute of Marine Science, Texas **5**: 251–264.
- MARCUS, ER. & MARCUS, EV. 1960. Opisthobranchs from American Atlantic warm waters. Bulletin of Marine Sciences of the Gulf and Caribbean **10**: 129–207.
- MARCUS, EV. & MARCUS, ER. 1962. Opisthobranchs from Florida and the Virgin Islands. Bulletin of Marine Sciences of the Gulf and Caribbean **12**: 450–488.
- MARCUS, EV. & MARCUS, ER. 1963. Opisthobranchs from the Lesser Antilles. Studies on the fauna of Curaçao and other Caribbean Islands **79**: 1–76.
- MARCUS, ER. & MARCUS, EV. 1967a. Tropical American Opisthobranch. Studies in Tropical Oceanography, Miami **6**: 1–137.
- MARCUS, EV. & MARCUS, ER. 1967b. Opisthobranchs from the Gulf of California. Studies in Tropical Oceanography, Institute of Marine Science **6**: 138–256.
- MARCUS, EV. & MARCUS, ER. 1967c. Opisthobranchs from the Southwestern Caribbean Sea. Bulletin of Marine Sciences **17** (3): 597–628.
- MARCUS, EV. & MARCUS, ER. 1968. *Flabellina engeli*, a new nudibranch from Curaçao. Beaufortia, Zoological Museum of the University of Amsterdam **15**: 139–142.
- MARCUS, ER. & MARCUS, EV. 1970. Opisthobranchs from Curaçao and faunistically related regions. Studies on the fauna of Curaçao and other Caribbean islands **33** (59): 1–129.
- MARCUS, EV. 1977. An annotated check list of the Western Atlantic warm water opisthobranchs. The Journal of the Molluscan Studies **Supplement 4**: 1–22.
- MARQUES, W. C.; MONTEIRO, I. O.; MÖLLER, O. O. & FERNANDES, E. H. 2006. Modelling the hydrodynamics of the Southern Brazilian Shelf. Proceedings of 8 ICSHMO, Foz do Iguaçu, Brazil, April 24–28. INPE 1311–1314.
- MUNIAIN C. 1997. Moluscos Opistobranquios de Argentina: Revisión Taxonómica y relación de Ecología Química en algunas especies patagónicas. Ph. D. Thesis, University of Oviedo, Spain.
- MUNIAIN, C. & ORTEA, J. 1999. First records of the genus *Berg-hia* Trinchese, 1877 (Opisthobranchia: Aeolidiidae) from Argentina, with description of a new species. Avicennia **10/11**: 143–150.
- ORTEA, J. 2001. El género Doto Oken, 1815 (Mollusca: Nudibranchia) en el mar Caribe: Historia Natural y descripción de nuevas especies. Avicennia **Supplement 3**: 1–46.
- ORTEA, J., CABALLER, M. & MORO, L. 2001. Una nueva especie del género *Noumea* Risbec, 1928 (Mollusca: Nudibranchia) del Caribe arrecifal de Costa Rica, descrita con motivo del Quinto Centenario de su descubrimiento. Avicennia **14**: 1–6.
- ORTEA, J. & ESPINOSA, J. 1996. Descripción de una nueva especie del género *Elysia* Risso, 1818 (Opisthobranchia: Sacoglossa) recolectada en Puerto Morelos, Mexico. Avicennia **4/5**: 115–119.
- ORTEA, J. & ESPINOSA, J. 1998. Estudio de nueve especies del género *Flabellina* Voight, 1834 (Mollusca: Nudibranchia) colectadas en Angola, Cabo Verde, Costa Rica, Cuba y Portugal, con la descripción de tres especies nuevas. Avicennia **8/9**: 135–148.
- ORTEA, J. & ESPINOSA, J. 2000. Nueva especie del género *Okeania* Menke, 1830 (Mollusca: Nudibranchia) de Cuba. Avicennia **12/13**: 84–86.
- ORTEA, J.; ESPINOSA, J. & CAMACHO, Y. 1999. Especies del género *Polycera* Cuvier, 1816 (Mollusca: Nudibranchia) recolectadas en la epifauna de algas rojas del Caribe de Costa Rica y Cuba. Avicennia **10/11**: 157–164.
- ORTEA, J. & MARTÍNEZ, E. 1997. Una nueva especie de *Chelidonura* A. Adams, 1850 (Mollusca: Opisthobranchia: Cephalaspidea) de las costas de Cuba. Avicennia **6/7**: 137–140.
- PADULA, V. & ABSALÃO, R. S. 2005. Primeiro Registro de *Babakina festiva* (Roller, 1972) (Mollusca: Nudibranchia) no Atlântico-Sul. Biociências, Porto Alegre **13**(2): 99–101.
- POLA, M.; CERVERA, J. L. & GOSLINER, T. M. 2005 A new species of *Tambja* Burn, 1962 (Nudibranchia, Polyceridae, Nembrothinae) from southern Brazil. Journal of the Marine Biological Association of the United Kingdom **85**: 979–984.
- REDFERN, C. 2001. Bahamian Seashells: a Thousand Species from Abaco, Bahamas. 280 pp.
- RIOS, E. 1994. Seashells of Brazil, 2nd Edition. Editora da FURG, Rio Grande, RS, Brazil. 368 pp.
- RIOS, E. C. & BARCELLOS, L. P. 1979. Nuevas ocurrencias de moluscos marinos para el Archipiélago de Fernando de Nor-

- onha, Brasil. Comunicaciones de la Sociedad Malacológica de Uruguay **5** (37): 163–166.
- ROCHA, L. A. 2003. Patterns of distribution and processes of speciation in Brazilian reef fishes. *Journal of Biogeography* **30**: 1161–1171.
- ROSENBERG, G. 2005. Malacolog 4.1: A Database of Western Atlantic Marine Mollusca. [WWW database (version 4.1.0)] URL <http://www.malacolog.org/>.
- SCHRÖDL, M. 1999. Zoogeographic relationships of the Magellanic Nudibranchia (Mollusca: Opisthobranchia) with particular reference to species from adjacent regions. *Scientia Marina* **63**(Supplement 1): 409–416.
- SNEATH, P. H. A. & SOKAL, R. R. 1973. Numerical Taxonomy. The principles and practice of numerical classification. WH Freeman, San Francisco.
- STORK, N. E. & SAMWAYS, M. J. 1995. Inventorying and Monitoring. Pp. 453–544 in: HEYWOOD, V.H. (ed.) *Global Biodiversity Assessment*. Cambridge University Press, Cambridge, UK.
- TEMPLADO, J.; LUQUE, A. A. & ORTEA, J. 1987. A new species of *Aegires* Löwen, 1844 (Opisthobranchia: Doridacea: Aegiridae) from the Caribbean Sea: *Aegires ortizi* sp. nov., with comparative descriptions of the North Atlantic species of this genus. *The Veliger* **29** (3): 303–307.
- THOMPSON, T. 1977. Jamaican Opisthobranch Molluscs I. *Journal of Molluscan Studies* **43**: 93–140.
- THOMPSON, T. E. 1980. Jamaican Opisthobranch Molluscs II. *Journal of Molluscan Studies* **46**: 74–99.
- TRONCOSO, J. S.; GARCÍA, F. J. & URGORRI, V. 1998. Anatomical data on rare *Hypselodoris picta* (Schultz, 1836) (Gastropoda, Doridacea) from the coast of Brazil with description of a new subspecies. *Bulletin of Marine Science* **63** (1): 133–141.
- VALDÉS, A.; HAMANN, J.; BEHRENS, D. W. & DUPONT, A. 2006. Caribbean seaslug. A field guide to the opisthobranch mollusks from the tropical northwestern Atlantic. *Sea Challengers Natural History Books, Etc.* Gig Harbor, Washington. Pp. 289.
- VERMEIJ, G. J. 1978. Biogeography and adaptation: Patterns of marine life. Harvard University Press, Cambridge.

Authors' addresses: Francisco J. GARCÍA (corresponding author), Departamento de Sistemas Físicos, Químicos y Naturales; Facultad Ciencias Experimentales; Universidad Pablo de Olavide, Sevilla, Spain, E-mail: fjgarcia@us.es; Marta DOMÍNGUEZ, Departamento de Ecología y Biología Animal, Facultad Ciencias del Mar, Universidad de Vigo, Campus de Vigo, E-36200, Vigo, Spain; Jesús S. TRONCOSO, Departamento de Ecología y Biología Animal, Facultad Ciencias del Mar, Universidad de Vigo, Campus de Vigo, E-36200, Vigo, Spain. E-mail: malvarez@uvigo.es; truncoso@uvigo.es.

Table 1. Species checklists and geographical distribution.

Species	CC-CR	CR-Cro	CRo-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Acteocina atrata</i> Mikkelsen, 1984	+	+	0	0	0	+	0	0	0	0	0	0	0	0
<i>Acteocina bernadensis</i> (Vanatta, 1901)	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Acteocina bidemata</i> (d'Orbigny, 1841)	+	+	+	+	0	+	0	+	+	+	+	+	+	+
<i>Acteocina bullata</i> (Kiener, 1834)	+	+	+	0	0	+	+	+	0	0	+	0	0	0
<i>Acteocina canaliculata</i> (Say, 1826)	+	+	+	0	0	0	0	0	0	0	0	0	0	0
<i>Acteocina candei</i> (d'Orbigny, 1841)	+	+	+	+	0	+	+	+	+	0	+	+	+	+
<i>Acteocina decurrens</i> (Verrill & Bush, 1900)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Acteocina inconspicua</i> Olsson & McGinty, 1958	+	+	+	0	+	+	0	0	0	0	0	0	0	0
<i>Acteocina kristensenii</i> Jong & Coomans, 1988	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Acteocina leptota</i> Woodring, 1928	+	+	+	0	0	+	+	0	0	0	0	0	0	0
<i>Acteocina liratispira</i> (E. A. Smith, 1872)	0	0	0	0	0	0	0	+	0	0	+	0	0	0
<i>Acteocina parvifica</i> (Dall, 1894)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Acteocina periplicata</i> (Dall, 1889)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Acteocina recta</i> (d'Orbigny, 1841)	+	+	+	0	0	+	0	0	0	+	0	0	0	0
<i>Acteon biplicatus</i> (Strebel, 1908)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Acteon candens</i> Rehder, 1939	+	+	+	0	0	+	0	0	0	0	+	0	0	0
<i>Acteon danaida</i> Dall, 1881	+	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Acteon delicatus</i> Dall, 1889	0	+	0	0	0	+	0	0	0	0	0	0	0	0
<i>Acteon exiguus</i> Möhrch, 1875	0	0	+	+	0	+	0	0	0	0	0	0	0	0
<i>Acteon fulvayi</i> McGinty, 1955	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Acteon incisus</i> Dall, 1881	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Acteon melanoides</i> Dall, 1881	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Acteon pelecus</i> Marcus, 1972	0	0	0	0	0	0	0	0	+	0	+	0	+	0
<i>Acteon perforatus</i> Dall, 1881	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Acteon splendidiolus</i> Möhrch, 1875	0	0	0	+	+	+	0	0	0	0	0	0	0	0
<i>Acteon vagabundus</i> (Mabille & Rochebrune, 1885)	0	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Aglaja felis</i> Marcus & Marcus, 1970	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Aglaja hummelincki</i> Marcus & Marcus, 1970	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Aglaja quitza</i> Marcus, 1979	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Aglaja unisa</i> Marcus & Marcus, 1969	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Alys alayoi</i> Espinosa & Ortea, 2004	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Alys caribaeus</i> (d'Orbigny, 1841)	+	+	+	0	0	+	0	0	+	0	0	0	0	0
<i>Alys guildingi</i> Sowerby II, 1869	0	0	+	0	0	+	0	0	+	0	0	0	0	0
<i>Alys macandrewii</i> E. A. Smith, 1872	0	+	+	0	+	+	0	0	+	0	0	0	0	0
<i>Alys riseanus</i> Möhrch, 1875	+	0	+	0	+	+	0	0	+	+	+	+	0	0
<i>Alys sandersoni</i> Dall, 1881	+	+	+	0	0	+	0	0	+	0	0	0	0	0
<i>Alys sharpi</i> Vanatta, 1901	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Bulla bernadensis</i> Verrill & Bush, 1900	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Bulla eburneola</i> (Dall, 1927)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Bulla krebssii</i> Dall, 1889	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Bulla nux</i> Menke, 1853	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Bulla persiciata</i> Menke, 1853	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Bulla pingicula</i> Seguenza, 1879	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Bulla solida</i> Gmelin, 1791	+	+	+	0	0	+	0	0	0	0	0	0	0	0

Species	CC-CR	CR-Cro	Cro-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Bulla striata</i> Bruguière, 1792	+	+	+	0	+	+	0	0	+	0	+	+	+	0
<i>Bullina exiguata</i> McGinny, 1955	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Bullina torrei</i> (Aguayo & Rehder, 1936)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Chelidomura boreolina</i> Marcus & Marcus, 1970	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Chelidomura cubana</i> Ortea & Martínez, 1997	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Chelidomura turpiniana</i> (Quoy & Gaimard, 1833)	+	0	0	0	+	0	0	0	0	0	0	0	0	0
<i>Chelidomura hummelinckii</i> Marcus & Marcus, 1970	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Chelidomura huancabosensis</i> Ortea & Espinosa, 1998	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Chelidomura muratoriae</i> Ortea, Espinosa & Moro, 2004	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Chelidomura petra</i> Marcus, 1976	0	0	0	0	0	+	0	0	+	0	0	0	0	0
<i>Chelidomura sabina</i> Marcus & Marcus, 1970	+	+	0	0	+	0	0	0	0	0	0	0	0	0
<i>Crenilabium exile</i> (Jeffreys, 1870)	+	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Cylindina alba</i> (Brown, 1827)	0	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Cylindina albertii</i> (d'Orbigny, 1841)	0	0	0	0	0	+	0	0	0	0	0	0	+	0
<i>Cylindina discus</i> Watson, 1883	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Cylindina elmeri</i> A. E. Verrill, 1885	+	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Cylindina georgiana</i> (Strebel, 1908)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Cylindina krebsii</i> Mörch, 1875	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Cylindina verrillii</i> Dall, 1889	+	0	0	0	0	0	0	0	0	0	0	0	+	0
<i>Cylindina vortex</i> (Dall, 1881)	+	+	0	0	+	0	0	+	+	0	0	0	0	0
<i>Cylindrobulla bevanii</i> P. Fischer, 1857	+	+	+	0	0	+	0	0	+	0	0	0	0	0
<i>Cylindrobulla gigas</i> Mikkelsen, 1998	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Diaphana caribaea</i> Espinosa, Ortea & Fernández-Garcés, 2001	0	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Diaphana seguenzæ</i> (Watson, 1886)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Gastropetron eliacmoi</i> Gosliner, 1989	+	+	0	0	0	+	0	0	0	0	0	+	0	0
<i>Gastropetron humanni</i> Gosliner, 1989	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Gastropetron rubrum</i> (Rafinesque, 1814)	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Gastropetron vesperillum</i> Gosliner & Armes, 1984	0	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Haminoea antillarum</i> (d'Orbigny, 1841)	+	+	0	0	0	+	0	0	+	0	0	0	0	0
<i>Haminoea elegans</i> (Gray, 1825)	+	+	+	0	+	+	0	0	+	+	0	+	+	0
<i>Haminoea glabra</i> (A. Adams, 1850)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Haminoea petiti</i> (d'Orbigny, 1841)	0	0	+	0	+	+	0	0	+	0	0	0	0	0
<i>Haminoea solitaria</i> (Say, 1822)	+	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Haminoea siccinca</i> (Conrad, 1846)	+	+	+	+	0	+	0	0	0	0	0	0	0	0
<i>Hydatina vesicaria</i> (Lightfoot, 1786)	+	+	+	+	0	+	0	0	+	+	0	0	0	0
<i>Laponacton pusillus</i> (Forbes, 1844)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Micromelo iudatus</i> (Bruguière, 1792)	+	0	+	0	0	+	0	0	+	+	0	0	0	0
<i>Myosoffia cumingii</i> (A. Adams, 1855)	+	0	0	0	0	0	0	0	0	0	+	+	+	0
<i>Myosoffia turrita</i> (Watson, 1883)	0	0	0	0	0	+	0	0	0	0	+	0	0	0
<i>Navanax acuminatus</i> (Bergh, 1893)	+	0	+	0	+	+	0	0	0	0	0	+	0	0
<i>Navanax orthynanus</i> (Rochebrune, 1881)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Onitacteon meeki</i> Dall, 1889	0	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Platinea alba</i> Mattox, 1958	0	0	+	0	0	0	0	0	0	0	0	0	+	0
<i>Platinea argentina</i> Carcelles, 1947	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Platinea caballeri</i> Ortea, Espinosa & Moro, 2001	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Platinea candiana</i> (d'Orbigny, 1841)	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Platinea falklandica</i> A. W. B. Powell, 1951	0	0	0	0	0	0	0	0	0	0	0	0	+	0

Species	CC-CR	CR-Cr ^o	CR-o-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Philine finmarchica</i> M. Sars, 1859	0	0	0	+	0	+	0	0	0	0	0	0	0	0
<i>Philine flexuosa</i> M. Sars, 1870	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Philine infundibulim</i> Dall, 1889	0	0	+	0	0	+	0	+	0	0	0	0	0	0
<i>Philine nera</i> Marcus & Marcus, 1969	0	0	0	0	0	0	0	0	0	0	+	+	+	0
<i>Philine planata</i> Dall, 1889	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Philine sagra</i> (d'Orbigny, 1841)	+	0	+	0	0	+	0	+	+	+	0	0	0	0
<i>Philine thurmanni</i> Marcus & Marcus, 1969	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Philine trachystraca</i> Watson, 1897	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Phlinopsis aeci</i> Ortea & Espinosa, 2001	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Phlinopsis pusa</i> (Marcus & Marcus, 1966)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Phocula euica</i> Marcus, 1953	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Pyrunculus caelatus</i> (Bush, 1885)	+	+	+	+	0	+	0	0	0	0	0	+	0	+
<i>Pyrunculus obesisculus</i> (Brugnone, 1877)	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Pyrunculus ovatus</i> (Jeffreys, 1871)	+	0	0	0	0	+	0	0	+	0	0	0	0	0
<i>Retusa canaliculata</i> Wells & Wells, 1962	0	0	0	0	0	0	0	0	+	0	0	+	0	0
<i>Retusa donita</i> (Dall, 1889)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Retusa frielei</i> (Dall, 1881)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Retusa onphalis</i> Mörch, 1875	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Retusa pervia</i> (Dall, 1889)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Retusa sosa</i> Marcus & Marcus, 1969	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Retusa spatha</i> (Watson, 1883)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Retusa sulcata</i> (d'Orbigny, 1841)	+	+	+	0	0	+	0	0	0	0	0	0	0	0
<i>Rictaxis punctostriatus</i> (C. B. Adams, 1840)	+	+	+	0	0	+	+	0	0	0	0	0	0	0
<i>Ringicula nitida</i> A. E. Verrill, 1872	0	+	0	0	0	+	+	0	0	0	0	0	0	0
<i>Ringicula semistriata</i> d'Orbigny, 1842	+	+	+	0	0	+	0	0	0	0	0	0	0	0
<i>Runcina divae</i> (Marcus & Marcus, 1963)	0	0	+	0	+	+	+	0	0	0	+	0	0	0
<i>Runcina inconspicua</i> A. E. Verrill, 1901	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Runcina prasina</i> (Mösch, 1863)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Scaphander bathynophilus</i> (Dall, 1881)	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Scaphander danius</i> Marcus & Marcus, 1967	0	0	+	0	0	+	0	+	0	+	0	0	0	0
<i>Scaphander loisae</i> Bullis, 1956	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Scaphander nobilis</i> A. E. Verrill, 1884	0	+	0	0	0	+	0	0	0	0	0	+	+	0
<i>Scaphander pilshryi</i> McGinty, 1955	0	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Scaphander punctostriatus</i> (Mighels & Adams, 1842)	0	+	+	0	0	+	0	0	0	0	0	0	0	0
<i>Scaphander watsoni</i> Dall, 1881	+	+	+	0	0	+	0	0	0	0	0	0	0	0
<i>Toledonia bullata</i> (Gould, 1847)	0	0	0	0	0	+	0	0	0	0	0	0	0	+
<i>Volvutella minuta</i> (Bush, 1885)	+	+	+	0	0	+	0	0	0	0	0	0	0	0
<i>Volvutella paupercula</i> (Watson, 1883)	0	+	0	0	0	+	0	0	0	0	0	0	0	0
<i>Volvutella persimilis</i> (Mösch, 1875)	+	+	+	+	0	+	+	0	+	0	0	+	+	0
<i>Volvutella recta</i> (Mösch, 1875)	0	0	0	0	0	+	0	0	0	0	+	0	0	0
<i>Volvutella texasiana</i> Harry, 1967	0	+	+	0	0	+	0	0	0	0	0	0	0	0
<i>Alderia modesta</i> (Loven, 1844)	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Aphysopsis formosa</i> Pruvot-Fol, 1953	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Ascobulla ulla</i> (Marcus & Marcus, 1970)	+	+	+	0	0	+	+	0	+	0	+	+	0	0
<i>Berthelina caribbea</i> Edmunds, 1963	+	0	+	0	0	+	0	0	+	0	0	0	0	0
<i>Bosellia coriinae</i> Marcus, 1973	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Bosellia marcusii</i> Marcus, 1972	+	0	+	0	0	+	0	0	0	0	0	0	0	0

Species	CC-CR	CR-Cro	Cro-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARB
<i>Bosellia mimetica</i> Trinchese, 1891	+	0	0	0	+	+	+	0	0	+	0	0	0	0
<i>Caliphylla mediterranea</i> A. Costa, 1867	+	0	0	0	+	+	0	0	+	0	0	+	0	0
<i>Costasiella lilianae</i> Marcus, 1969	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Costasiella nomatoi</i> Marcus & Marcus, 1960	+	0	0	0	0	+	0	0	0	0	0	+	0	0
<i>Costasiella ocellifera</i> (Simroth, 1895)	+	0	+	0	0	+	+	0	0	0	0	+	0	0
<i>Cyerce unillensis</i> Engel, 1927	+	0	0	0	+	+	+	0	0	0	0	0	0	0
<i>Cyerce cristallina</i> (Trinchese, 1881)	+	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Cyerce edmundsi</i> Thompson, 1977	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Cyerce habonensis</i> Ortea & Templado, 1989	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Elysia canguinua</i> Marcus, 1955	+	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Elysia carinus</i> (Gould, 1870)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Elysia carze</i> Marcus, 1957	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Elysia chinui</i> Marcus, 1955	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Elysia chlorotica</i> Gould, 1870	+	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Elysia cornigera</i> Nuttall, 1889	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Elysia eugeniiae</i> Ortea & Espinosa, 2002	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Elysia evrhuae</i> Marcus, 1957	+	0	+	0	0	0	0	0	0	0	0	+	0	0
<i>Elysia flavo</i> A. E. Verrill, 1901	0	0	+	0	0	+	+	0	+	0	0	0	0	0
<i>Elysia misbeti</i> Thompson, 1977	0	0	0	0	0	+	+	0	0	0	0	0	0	0
<i>Elysia ornata</i> (Swainson, 1840)	+	0	+	0	+	+	+	0	+	0	0	0	0	0
<i>Elysia papillosa</i> A. E. Verrill, 1901	+	0	+	0	+	+	+	0	0	0	0	0	0	0
<i>Elysia patagonica</i> Munnin & Ortea, 1997	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Elysia patina</i> Marcus, 1980	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Elysia pratensis</i> Ortea & Espinosa, 1996	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Elysia punctonii</i> Thompson, 1977	0	0	+	0	+	0	0	0	0	0	0	0	0	0
<i>Elysia serca</i> Marcus, 1955	+	+	0	0	+	+	0	0	0	0	0	+	0	0
<i>Elysia subornata</i> A. E. Verrill, 1901	+	0	0	0	+	+	+	0	0	0	0	+	0	0
<i>Elysia tica</i> Marcus & Marcus, 1967	+	0	+	0	+	+	+	0	+	+	0	0	0	0
<i>Elysia zutetiae</i> Ortea & Espinosa, 2002	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Ercolanina circetus</i> (Marcus & Marcus, 1970)	+	0	+	0	+	+	+	0	0	0	0	0	0	0
<i>Ercolanina fumerea</i> (A. Costa, 1867)	+	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Ercolanina fuscata</i> (Gould, 1870)	+	0	0	0	+	+	0	0	0	0	0	+	0	0
<i>Ercolanina selva</i> Ortea & Espinosa, 2001	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Hermatea bifida</i> (Montagu, 1815)	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Hermatea corrida</i> Marcus, 1955	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Hermatea cruciata</i> Gould, 1870	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Lobiger souverbii</i> Fischer, 1857	+	+	+	0	+	+	0	0	0	0	0	+	0	0
<i>Moungonia germaniae</i> Marcus & Marcus, 1970	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Moungonia mirra</i> Marcus & Marcus, 1970	0	0	0	0	+	0	0	0	0	0	0	0	0	0
<i>Oxytne agurroi</i> Jaume, 1945	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Oxytne antillarum</i> Mörch, 1863	+	+	+	0	+	+	+	0	+	+	+	+	0	0
<i>Oxytne azuopunctata</i> Jensen, 1980	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Oxytne panamensis</i> Pilsbry & Olsson, 1943	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Phyllabrac hillius viridis</i> (Deshayes, 1857)	+	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Placida dendritica</i> (Alder & Hancock, 1843)	+	0	0	0	+	0	0	0	0	0	0	0	0	0
<i>Placida kingsoni</i> Thompson, 1977	+	+	+	0	+	+	0	0	0	0	0	0	0	0
<i>Placida verticillata</i> Ortea, 1981	+	0	+	0	0	0	0	0	0	0	0	0	0	0

Species	CC-CR	CR-Cr ^o	CR ^o -O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Stiliger fuscovittatus</i> Lance, 1962	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Stiliger talis</i> Marcus, 1856	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Stiliger vanellus</i> (Marcus, 1957)	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Stiliger vossi</i> Marcus & Marcus, 1960	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Thuridilla mazda</i> Ortea & Espinosa, 2000	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Thuridilla picta</i> (A. E. Verrill, 1901)	0	0	0	0	0	+	+	0	0	0	0	0	0	0
<i>Tridachia crispata</i> (Mörch, 1863)	+	+	+	0	+	+	+	0	0	0	0	0	0	0
<i>Tridachia sclerannii</i> (Mörch, 1863)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Vohvarella bernudae</i> Clark, 1982	+	0	+	0	0	0	+	0	0	0	0	0	0	0
<i>Akera bayeri</i> Marcus & Marcus, 1967	+	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Aphysia cervina</i> (Dall & Simpson, 1901)	+	0	+	0	0	+	0	0	+	0	0	+	0	0
<i>Aphysia dactylorella</i> Rang, 1828	+	+	+	0	+	+	+	0	+	+	+	+	0	0
<i>Aphysia fasciata</i> Poiret, 1789	+	+	+	0	+	+	+	0	0	0	+	+	0	0
<i>Aphysia geographica</i> (Adams & Reeve, 1850)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aphysia juliana</i> Quoy & Gaimard, 1832	+	0	0	0	+	+	0	0	+	0	0	+	0	0
<i>Aphysia morio</i> (Verrill, 1901)	+	+	0	0	0	0	+	0	0	0	0	0	0	0
<i>Aphysia parvula</i> Mörch, 1863	0	+	+	0	0	0	+	0	+	+	+	0	0	0
<i>Aphysia willcoxi</i> Heilprin, 1887	+	+	0	0	0	+	+	0	0	0	0	0	0	0
<i>Bursatella leachii</i> Blainville, 1817	+	+	+	0	+	+	+	0	+	+	+	+	+	0
<i>Dolabrifera dolabrifera</i> (Rang, 1828)	+	0	+	+	+	+	+	0	0	0	+	0	0	0
<i>Notarchus punctatus</i> Philippi, 1836	0	0	0	0	0	+	0	0	0	+	0	0	0	0
<i>Petalifera petalifera</i> (Rang, 1828)	0	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Petalifera ramosa</i> Baba, 1959	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Phyllaphysia engelii</i> Marcus, 1955	+	+	+	0	0	+	0	0	+	+	0	+	0	0
<i>Phyllaphysia smaragda</i> Clark, 1977	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Sylocheilus citrinus</i> (Rang, 1828)	0	0	0	0	0	0	0	0	+	0	0	+	0	0
<i>Sylocheilus longicauda</i> (Quoy & Gaimard, 1824)	0	0	0	0	0	+	0	0	+	0	0	+	0	0
<i>Sylocheilus striatus</i> (Quoy & Gaimard, 1832)	+	0	+	0	+	+	+	0	+	+	+	0	0	0
<i>Berthella agassizii</i> (MacFarland, 1909)	0	0	+	0	+	0	+	0	+	0	+	+	0	0
<i>Berthella patagonica</i> (d'Orbigny, 1835)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Berthella stellata</i> (Risso, 1826)	+	0	+	0	0	+	0	0	+	0	+	+	0	0
<i>Berthella taitii</i> Marcus, 1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Berthellina circularis</i> (Mörch, 1863)	0	0	0	0	0	+	0	0	+	0	0	0	0	0
<i>Berthellina quadridens</i> (Mörch, 1863)	0	0	0	0	0	+	0	0	+	+	0	0	0	0
<i>Pleurobranchaea agassizii</i> Bergh, 1897	+	0	0	0	0	+	0	0	0	+	0	0	0	0
<i>Pleurobranchaea inconspicua</i> Bergh, 1897	+	+	+	+	0	+	0	0	+	+	+	+	+	+
<i>Pleurobranchaea obesa</i> (Verrill, 1882)	0	+	0	0	0	+	0	0	0	0	0	0	0	0
<i>Pleurobranchaea tarda</i> Verrill, 1880	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Pleurobranchaea areolata</i> Mörch, 1863	+	+	+	+	+	+	+	0	+	0	0	+	0	0
<i>Pleurobranchus enysii</i> Marcus, 1984	0	0	+	0	0	+	0	0	+	0	0	0	0	0
<i>Pleurobranchus evelinae</i> Thompson, 1977	0	0	0	0	0	+	0	0	+	0	0	0	0	0
<i>Pleurobranchus ionipsi</i> Marcus, 1984	0	0	0	0	0	0	0	0	+	0	0	+	0	0
<i>Pleurobranchus lacteus</i> Dall & Simpson, 1901	0	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Pleurobranchus niveus</i> (Verrill, 1901)	0	0	0	0	0	+	+	0	0	0	0	0	0	0
<i>Tylodina americana</i> Dall, 1890	0	+	0	0	0	+	+	0	0	0	0	0	0	0
<i>Umbraculum umbraculum</i> (Lightfoot, 1786)	+	+	+	0	0	+	+	+	+	+	0	0	0	0
<i>Aegires absaloi</i> García, Troncoso & Dominguez, 2002	0	0	0	0	0	0	0	0	+	0	0	0	0	0

Species	CC-CR	CR-Cro	Cro-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Aegires gonaezi</i> Ortea, Luque & Templado, 1990	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Aegires ortzi</i> Templado, Luque & Ortea, 1987	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Aegires sublaevis</i> Odhner, 1932	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Aicula espinosai</i> Ortea, 2001	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Aphelodoris antilleensis</i> Bergh, 1879	+	0	+	0	+	+	0	0	0	0	0	0	0	0
<i>Aporodoris millegrana</i> (Alder & Hancock, 1854)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Atagema browni</i> Thompson, 1980	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Atagema hispidula</i> (d'Orbigny, 1837)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Atagema preta</i> (Marcus & Marcus, 1967)	+	0	0	0	0	+	0	0	0	0	+	0	0	0
<i>Cadlina rimaia</i> Marcus, 1955	+	0	+	0	+	0	0	0	0	0	0	0	0	0
<i>Cadlina scabriuscula</i> (Bergh, 1890)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Cadlina sparsa</i> (Odhner, 1921)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Ceratophyllidia papilligera</i> (Bergh, 1890)	0	+	+	0	+	+	+	+	+	0	0	0	0	0
<i>Chironodoris binza</i> Marcus & Marcus, 1963	0	0	0	0	0	+	0	0	0	0	+	0	0	0
<i>Chironodoris clientia</i> (Russell, 1935)	+	+	+	0	0	+	0	0	+	0	0	+	0	0
<i>Chironodoris dieyi</i> Marcus & Marcus, 1970	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Chironodoris gyalumii</i> Thompson, 1980	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Chironodoris kempfi</i> Marcus, 1971	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Chironodoris neona</i> (Marcus, 1955)	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Chironodoris perla</i> Marcus, 1976	0	0	+	0	+	0	0	0	0	0	0	0	0	0
<i>Chironodoris poriga</i> Marcus & Marcus, 1970	0	0	0	0	+	0	0	0	0	0	0	0	0	0
<i>Chironodoris punctilucens</i> Bergh, 1890	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Chironodoris roseopicta</i> Verrill, 1900	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Corambe exelinae</i> Marcus, 1958	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Corambe obscura</i> A. E. Verrill, 1870	0	+	0	0	0	+	0	0	0	0	0	0	0	0
<i>Dendrodois brepsii</i> (Mölich, 1863)	+	+	+	0	+	+	0	0	0	0	+	0	0	0
<i>Dendrodois magagnai</i> Ortea & Espinosa, 2001	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Dendrodois seNEGALensis</i> Bouchet 1977	0	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Dendrodois warra</i> Marcus & Gallagher, 1976	+	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Dianthula greeleyi</i> (MacFarland, 1909)	0	0	+	0	0	+	0	0	+	0	+	0	0	0
<i>Dianthula punctulata</i> (d'Orbigny, 1837)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Dianthula worki</i> (Marcus & Marcus, 1967)	+	0	0	0	+	0	0	0	0	0	0	0	0	0
<i>Discodoris alba</i> White, 1952	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Discodoris branneri</i> MacFarland, 1909	0	0	0	0	0	0	0	0	+	0	0	0	0	0
<i>Discodoris exelinae</i> Marcus, 1955	+	+	+	0	+	+	0	0	0	0	+	0	0	0
<i>Discodoris morrisensis</i> Marcus & Marcus, 1963	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Discodoris mitta</i> Bergh, 1877	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Discodoris nothia</i> Bergh, 1877	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Discodoris ketos gila</i> (Marcus & Marcus, 1970)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Discodoris phoca</i> Marcus & Marcus, 1967	+	0	+	0	0	0	0	0	+	0	0	0	0	0
<i>Discodoris purcina</i> Marcus & Marcus, 1967	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Discodoris vonthierigi</i> MacFarland, 1909	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Doridella caranbola</i> (Marcus, 1955)	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Doridopsilla areolata nigrolucata</i> Meyer, 1977	0	0	+	0	0	0	0	0	0	0	0	+	0	0
<i>Doridopsilla espinosai</i> Valdés & Ortea, 1998	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Doridopsilla pharpha</i> Marcus, 1961	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Doris bicolor</i> (Bergh, 1884)	+	0	0	0	0	0	0	0	0	0	0	0	0	0

Species	CC-CR	CR-Cr ⁰	CR-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Doris boveana</i> Marcus, 1955	0	0	0	0	+	0	0	0	0	0	0	+	0	0
<i>Doris fontainei</i> d'Orbigny, 1837	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Doris fretterae</i> Thompson, 1980	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Doris ilo</i> (Marcus, 1955)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Doris kerguelensis</i> (Bergh, 1884)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Doris kyolis</i> (Marcus & Marcus, 1967)	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Doris pickensi</i> Marcus & Marcus, 1967	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Doris verrucosa</i> Linné, 1758	+	+	+	0	0	0	0	0	+	0	+	+	0	0
<i>Etidoris ladiavii</i> Ihering, 1886	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Galacera marplatensis</i> (Franceschi, 1928)	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Garganella iminiaculata</i> Bergh, 1894	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Geitodoris patagonica</i> Odhner, 1926	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Geitodoris planata</i> (Alder & Hancock, 1846)	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Geitodoris pusae</i> (Marcus, 1955)	+	0	0	0	+	+	0	+	0	0	0	+	0	+
<i>Geitodoris immunda</i> Bergh, 1894	0	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Glossodoris moerchi</i> (Bergh, 1879)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Glossodoris sedna</i> (Marcus & Marcus, 1967)	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Goniodoris mimula</i> Marcus, 1955	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Hallaxa apetae</i> Marcus, 1957	0	0	0	0	0	0	0	0	0	0	+	+	0	0
<i>Hexabranchius noronensis</i> Marcus & Marcus, 1962	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Hopllocampus pupposus</i> Odhner, 1926	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Hoplodoris hiansosorum</i> Dominguez, Garcia & Troncoso, 2006	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Hypselodoris aciriba</i> Marcus & Marcus, 1967	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Hypselodoris bayeri</i> (Marcus & Marcus, 1967)	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Hypselodoris espinosai</i> Ortea & Valdés, 1996	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Hypselodoris lajensis</i> Troncoso, Garcia & Urgorri, 1998	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Hypselodoris marci</i> Marcus, 1971	0	0	+	0	0	+	0	0	0	0	+	0	0	0
<i>Hypselodoris picta</i> (Schultz, 1836)	+	+	0	0	0	0	0	0	0	0	0	+	0	0
<i>Hypselodoris ruthae</i> Marcus & Hughes, 1974	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Hypselodoris sycilla</i> (Bergh, 1890)	+	0	0	0	0	+	0	0	+	0	0	0	0	0
<i>Hypselodoris zebra</i> (Heilprin, 1888)	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Jorunna spazzola</i> (Marcus, 1955)	0	0	0	0	0	+	0	0	0	+	+	+	0	0
<i>Kankelibranchius incognitus</i> Ortea, Espinosa & Caballer, 2005	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Lophodoris scala</i> Marcus & Marcus, 1970	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Mexichronis mollot</i> Ortea & Valdés, 1996	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Noumea regalis</i> Ortea, Caballer & Moro, 2001	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Okenia impexa</i> Marcus, 1957	0	0	0	0	0	+	0	0	0	0	0	+	0	0
<i>Okenia miramarae</i> Ortea & Espinosa, 2000	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Okenia zoobotryon</i> (Smallwood, 1910)	+	0	0	0	0	+	+	0	0	0	+	+	0	0
<i>Onchidoris aureopuncta</i> (Verrill, 1901)	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Onchidoris bilamellata</i> (Linné, 1761)	0	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Onchidoris lactea</i> (Verrill, 1900)	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Onchidoris miniata</i> (Verrill, 1901)	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Onchidoris olivacea</i> (Verrill, 1900)	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Onchidoris quadrimaculata</i> (Verrill, 1900)	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Paradoris mulciber</i> (Marcus, 1971)	0	0	+	0	0	0	0	0	+	0	0	0	0	0
<i>Peltodoris crucei</i> (Mösch, 1863)	0	0	0	0	0	+	0	0	0	0	0	0	0	0

Species	CC-CR	CR-Cn	CR-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Petioloris hummelinckii</i> Marcus & Marcus, 1963	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Platythelidia molaeusis</i> (Meyer, 1977)	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Platyloris angustipes</i> (Mölich, 1863)	+	0	+	0	0	+	0	0	+	0	0	0	0	0
<i>Placompherus gulo</i> Marcus, 1979	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Placompherus lucayensis</i> Hamann & Farmer, 1988	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Placompherus pilatensis</i> Hamann & Farmer, 1988	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Polycera antistila</i> Marcus, 1957	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Polycera hedgpethi</i> Marcus, 1964	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Polycera herthae</i> Marcus & Marcus, 1963	0	0	0	0	+	0	0	0	0	0	0	0	0	0
<i>Polycera hummi</i> Abbott, 1952	+	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Polycera japonica</i> Baba, 1949	0	0	0	0	0	+	0	0	0	0	0	+	0	0
<i>Polycera manzanilloensis</i> Ortea, Espinosa & Camacho, 1999	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Polycera odlineri</i> Marcus, 1955	+	0	0	0	0	+	0	0	0	0	0	+	0	0
<i>Polycera quadrilineata</i> (Müller, 1776)	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Polycera rycia</i> Marcus & Marcus, 1970	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Polycerella emertoni</i> Verrill, 1880	+	0	+	0	0	+	0	0	0	0	0	+	0	0
<i>Risbecia nydya</i> (Marcus & Marcus, 1967)	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Rostanga hydra</i> Marcus, 1958	0	0	0	0	0	0	0	0	+	0	0	0	0	+
<i>Rostanga pulchra</i> MacFarland, 1905	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Siratus ilo</i> Marcus, 1955	0	0	0	0	0	0	0	0	0	+	+	0	0	0
<i>Siratus kvolds</i> Marcus & Marcus, 1967	0	0	0	0	0	0	0	0	+	0	+	0	0	0
<i>Tamblya divae</i> (Marcus, 1958)	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Tamblya gratiosa</i> (Bergh, 1890)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Tamblya oliva</i> Meyer, 1977	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Tamblya stegosauriformis</i> Pola, Cervera & Gosliner, 2005	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Taranga millegrana</i> (Alder & Hancock, 1855)	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Taranga telopia</i> Marcus, 1955	+	0	+	0	0	+	0	0	0	0	+	0	0	0
<i>Taranga tritorquis</i> Ortea, Pérez & Llera, 1982	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Thecacera pennigera</i> (Montagu, 1815)	+	0	0	0	0	0	0	0	+	0	0	+	0	0
<i>Thorlisa dhinda</i> Marcus, 1955	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Thorlisa lutea</i> (Marcus & Marcus, 1967)	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Trapania dubya</i> Marcus, 1972	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Trapania maringa</i> Marcus, 1957	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Trippa anceps</i> (Bergh, 1890)	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Tyrinna evelinae</i> (Marcus, 1958)	0	0	0	0	0	+	0	0	0	0	+	0	0	0

Species	CC-CR	CR-CrO	CRo-O	O-A	ABC	WI	BER	AS	NBS	EBS	ACR	SBB	SBS	ARG
<i>Doto pita</i> Marcus, 1955	+	0	0	0	+	+	0	0	0	0	0	+	0	0
<i>Doto prorata</i> Ortea, 2001	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Doto pygmaea</i> Bergh, 1871	0	0	0	0	0	+	+	0	0	0	0	0	0	0
<i>Doto sabuli</i> Ortea, 2001	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Doto tiva</i> Marcus, 1955	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Doto varaderoensis</i> Ortea, 2001	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Hancockia ryrca</i> Marcus, 1957	0	0	0	0	+	+	0	0	0	0	0	+	0	0
<i>Lomanotus phiops</i> Marcus, 1957	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Lomanotus verniformis</i> Eliot, 1908	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Mariontia cucullata</i> (Gould, 1852)	0	0	+	0	0	0	0	0	0	0	0	+	0	+
<i>Mariontia tedi</i> Marcus, 1983	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Miesea evelinae</i> (Marcus, 1957)	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Scyllaea pelagica</i> Linné, 1758	+	+	+	0	+	+	+	0	+	0	0	0	0	0
<i>Phylliroe atlantica</i> Bergh, 1871	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Phylliroe bucephala</i> Lamarck, 1816	0	0	0	0	0	0	+	0	0	0	0	0	0	0
<i>Tethys occidentalis</i> (Odmer, 1936)	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Tritonia australis</i> Bergh, 1898	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Tritonia bayeri</i> Marcus & Marcus, 1967	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Tritonia ertosi</i> Marcus, 1983	0	0	0	0	0	0	0	0	0	0	0	0	+	+
<i>Tritonia launerorum</i> Gosliner & Ghiselin, 1987	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Tritonidoxa wellsi</i> (Marcus, 1961)	+	0	+	0	0	0	0	0	0	0	0	+	0	0
<i>Tritoniopsis frydis</i> Marcus & Marcus, 1970	+	0	0	0	0	+	+	0	0	0	0	0	0	0
<i>Arminia elongata</i> Ardila & Valdés, 2004	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Arminia juliana</i> Ardila & Diaz, 2002	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Arminia muelleri</i> (Ihering, 1886)	+	+	+	+	0	+	0	0	0	0	+	+	0	0
<i>Janolus comis</i> Marcus, 1955	+	0	0	0	+	0	0	0	0	0	0	+	0	0
<i>Janolus costacubensis</i> Ortea & Espinosa, 2000	0	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Janolus micloc</i> (Marcus, 1958)	+	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Anteaeolidiella indica</i> Bergh, 1888	+	0	0	0	+	+	0	0	0	0	+	+	0	+
<i>Aeolidia serotina</i> Bergh, 1873	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aeolidiella occidentalis</i> Bergh, 1875	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Anetarca brasiliana</i> García & Troncoso, 2004	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Austracolis catina</i> Marcus & Marcus, 1967	+	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Babakina festiva</i> (Roller, 1972)	0	0	0	0	0	0	0	0	0	0	+	0	0	0
<i>Berghia bentvea</i> (Marcus, 1958)	0	0	0	0	0	0	0	0	0	0	+	+	0	0
<i>Berghia coerulescens</i> (Laurillard in Cuvier, 1830)	0	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Berghia creutzbergi</i> Marcus & Marcus, 1970	0	0	+	0	+	+	0	0	0	0	+	0	0	0
<i>Berghia verrucicornis</i> (A. Costa, 1864)	+	0	+	0	0	+	0	0	0	0	0	0	0	0
<i>Berghia rissodominguezi</i> Muniaín & Ortea, 1999	0	0	0	0	0	0	0	0	0	0	0	0	0	+
<i>Calinella bandelli</i> Marcus, 1976	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Catironia nana</i> Marcus & Marcus, 1960	+	0	0	0	+	+	0	0	0	0	0	0	0	0
<i>Catironia oba</i> Marcus, 1970	0	0	0	0	0	0	0	0	0	0	0	+	0	0
<i>Cerberilla tanna</i> Marcus & Marcus, 1960	0	+	0	0	0	0	0	0	0	0	0	0	0	0
<i>Cratena pilata</i> (Gould, 1870)	+	+	0	0	0	0	+	0	0	0	0	+	0	0
<i>Cratena pituitensis</i> Ortea, Caballer & Espinosa, 2003	0	0	+	0	0	0	0	0	0	0	0	0	0	0
<i>Cuthona barbadiana</i> Edmunds & Just, 1983	0	0	0	0	0	+	0	0	0	0	0	0	0	0
<i>Cuthona coerulea</i> (Montagu, 1804)	+	0	0	0	0	0	0	0	0	0	+	+	0	0

[illegible]

Table 2. Limits and features of the areas considered along Brazilian coasts, based on CASTRO & MIRANDA (1998).

Regions	Coastal Limits	Width of continental shelf (Km)	Shelf-breaks depth (m)	Currents	Salinity (‰)	Temperature (°C)
Amazon Shelf (AS)	4°N–2°S	125–320	140	North Brazil Current Geostrophic Current Amazon River	<33	27
Northeastern Brazilian Shelf (NBS)	2°S–8°S	40–85	73	North Brazil Current	36–37	Summer: 27–29 Winter: 26–28
Eastern Brazilian Shelf (EBS)	8°S–15°S	10–15	50–60	South Equatorial Current	36–37 River 32–33	Summer: 27–28 Winter: 25–26
Abrolhos-Campos region (ACR)	15°S–23°S	35–190	60–100	Brazil Current South Atlantic Central Water Coastal Water Upwelling events	36.5–37	Summer: 25–27 Winter: 22–24 Upwelling: 16
South Brazilian Bight (SBB)	23°S–28.4°S	50–230	120–180	Brazil Current South Atlantic Central Water Fresh Coastal Water	<33–>36	Summer: 25–27 Upwelling in North: 21 Winter: 20–23 Water from South <18
Southern Brazilian Shelf (SBS)	28.5°S–34°S	110–170	180	Summer: Brazil Current Winter: Subantarctic Water	Summer: >36 Winter: <34	Summer: >20 Winter: <15

Table 3. Number of species by order or suborder for each area considered.

	CC-CR	CR-Cro	Cro-O	ABC	WI	BER	O-A	AS	NBS	EBS	ACR	SBB	SBS	Arg
Cephalaspidea	50	39	52	14	88	19	8	9	26	11	14	19	14	10
Sacoglossa	38	8	29	19	37	14	0	0	7	3	3	18	0	1
Anaspidea	14	7	11	10	14	8	1	0	10	6	6	8	1	0
Notaspidea	6	5	8	3	12	5	1	1	10	4	3	5	1	2
Doridina	35	12	39	13	52	11	0	2	17	2	23	36	0	14
Dendronotina	9	2	16	6	14	5	1	0	3	1	0	8	1	3
Arminina	3	1	4	1	2	0	1	0	0	0	1	2	0	0
Aeolidina	18	5	19	19	31	10	0	0	2	1	12	26	1	5
Total	173	79	178	85	250	72	12	12	75	28	62	122	18	35

Table 4. Percentage of species from each area (vertical column) present in other localities.

	CC-CR	CR-Cro	Cro-O	ABC	WI	Ber	O-A	AS	NBS	EBS	ABR	SBB	SBS	ARG	Total of species
CC-CR	100	34	53	34	73	25	5	4	24	11	19	34	6	3	173
CR-Cro	75	100	66	30	78	39	9	8	34	15	25	37	13	8	79
Cro-O	51	29	100	24	69	21	6	5	28	12	20	25	4	3	178
ABC	69	28	51	100	80	34	2	2	27	18	28	44	5	2	85
WI	50	25	49	27	100	19	5	4	21	10	16	24	4	2	250
BER	61	43	51	40	67	100	6	6	32	17	26	26	8	4	72
O-A	67	58	83	17	100	33	100	17	50	25	42	50	33	4	42
AS	58	50	75	17	75	33	17	100	58	42	42	33	25	25	12
NBS	55	36	65	31	69	31	8	9	100	28	31	39	12	7	75
EBS	68	43	75	54	86	43	11	18	75	100	46	39	18	7	28
ABR	53	32	58	39	65	31	8	8	37	21	100	61	16	5	62
SBB	48	24	36	30	48	16	5	3	24	9	31	100	10	7	122
SBS	56	56	44	22	61	33	22	17	50	28	56	67	100	28	18
ARG	17	17	17	6	17	9	14	9	14	6	9	26	14	100	35
Total of species	173	79	178	85	250	72	12	12	75	28	62	122	18	35	

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Bonn zoological Bulletin - früher Bonner Zoologische Beiträge.](#)

Jahr/Year: 2007

Band/Volume: [55](#)

Autor(en)/Author(s): García Francisco J., Domínguez Marta, Troncoso Jesús S.

Artikel/Article: [Biogeographic considerations of the Opisthobranchia \(Mollusca: Gastropoda\) fauna from the Brazilian littoral and nearby areas 203-222](#)